

BP-2010E/98E References

1. Li, M., et al., Baicalin mitigates hypertension-linked alterations in the intestinal lymphatic vasculature in part through preserving the functional barrier integrity of lymphatic endothelial cells. *Biomedicine & Pharmacotherapy*, 2023. 160: p. 114418.
2. Cai, X., et al., Noninvasive Stereotactic Radiotherapy for Renal Denervation in a Swine Model. *Journal of the American College of Cardiology*, 2019. 74(13): p. 1697-1709.
3. Watanabe, H., et al., Protective Effects of Japanese Soybean Paste (Miso) on Stroke in Stroke-Prone Spontaneously Hypertensive Rats (SHRSP). *Am J Hypertens*, 2017. 31(1): p. 43-47.
4. Chen, W., et al., Effects and Mechanisms of Radiofrequency Ablation of Renal Sympathetic Nerve on Anti-Hypertension in Canine. *Arq Bras Cardiol*, 2017. 108(3): p. 237-245.
5. Yamashita, M., et al., Podocyte-specific NF-kappaB inhibition ameliorates proteinuria in adriamycin-induced nephropathy in mice. *Clin Exp Nephrol*, 2016. 2016: p. 18.
6. Nakano, S., et al., The effect of chronic renal failure on cardiac function: an experimental study with a rat model. *J Echocardiogr*, 2016. 2016: p. 14.
7. Matsubara, K., et al., Immune activation during the implantation phase causes preeclampsia-like symptoms via the CD40-CD40 ligand pathway in pregnant mice. *Hypertens Res*, 2016.
8. Li, D., et al., A Novel Swine Model of Spontaneous Hypertension With Sympathetic Hyperactivity Responds Well to Renal Denervation. *American Journal of Hypertension*, 2015.
9. Kitamura, A., et al., Gradual cerebral hypoperfusion in spontaneously hypertensive rats induces slowly evolving white matter abnormalities and impairs working memory. *J Cereb Blood Flow Metab*, 2015. 2015(13).
10. Yoshida, T., et al., Kruppel-like factor 4 protein regulates isoproterenol-induced cardiac hypertrophy by modulating myocardin expression and activity. *J Biol Chem*, 2014. 289(38): p. 26107-18.
11. Yoshida, T., et al., Deletion of Kruppel-like factor 4 in endothelial and hematopoietic cells enhances neointimal formation following vascular injury. *J Am Heart Assoc*, 2014. 3(1): p. 000622.
12. Ma, L.Q., et al., Post-treatment with Ma-Xing-Shi-Gan-Tang, a Chinese medicine formula, ameliorates lipopolysaccharide-induced lung microvessel hyperpermeability and inflammatory reaction in rat. *Microcirculation*, 2014. 2014(8): p. 12144.
13. Yoshida, T., et al., Smooth Muscle-Selective Inhibition of Nuclear Factor-kappaB Attenuates Smooth Muscle Phenotypic Switching and Neointima Formation Following Vascular Injury. *J Am Heart Assoc*, 2013. 2(3): p. 000230.
14. Sun, W., et al., [Anaphylactoid reactions induced by polysorbate 80 on Beagle dogs]. *Zhongguo Zhong Yao Za Zhi*, 2012. 36(14): p. 1874-8.
15. 王志国, et al., 组胺致 Beagle 犬类过敏反应实验模型研究. *中国中药杂志*, 2011(14): p. 1842-1844.
16. 孙伟伟, et al., 多种聚氧乙烯脱水山梨醇单油酸酯致 Beagle 犬类过敏反应的试验研究. *中国中药杂志*, 2011(14): p. 1874-1878.
17. Yatabe, J., et al., Angiotensin III stimulates aldosterone secretion from adrenal gland partially via angiotensin II type 2 receptor but not angiotensin II type 1 receptor. *Endocrinology*, 2011. 152(4): p. 1582-8.
18. Sugiyama, T., et al., Effects of fasudil, a Rho-associated protein kinase inhibitor, on optic nerve head blood flow in rabbits. *Invest Ophthalmol Vis Sci*, 2011. 52(1): p. 64-9.
19. Shibata, M., et al., Changes in optic nerve head blood flow, visual function, and retinal histology in hypercholesterolemic rabbits. *Exp Eye Res*, 2011. 93(6): p. 818-24.
20. Shibata, M., et al., Disruption of gap junctions may be involved in impairment of autoregulation in optic nerve head blood flow of diabetic rabbits. *Invest Ophthalmol Vis Sci*, 2011. 52(5): p. 2153-9.
21. Kusumoto, K., et al., Antihypertensive, insulin-sensitising and renoprotective effects of a novel, potent and long-acting angiotensin II type 1 receptor blocker, azilsartan medoxomil, in rat and dog models. *Eur J Pharmacol*, 2011. 669(1-3): p. 84-93.
22. Hu, X.-B., et al., Intravascular ultrasound area strain imaging used to characterize tissue components and assess vulnerability of atherosclerotic plaques in a rabbit model. *Ultrasound in Medicine & Biology*, 2011. 37(10): p.

23. 张美玉, 静脉用鱼腥草注射剂致敏性评价和比较药理学研究——博士论文. 2010, 中国中医科学院.
24. 卢祺炯, et al., 二种方法监测麻醉状态下 Beagle 犬血压动态变化的比较. 科技创新导报, 2010(29): p. 255-256.
25. Xin, Y.F., et al., Stereospecific pharmacodynamics and chiral inversion of NG-nitro-arginine in the beagle dogs. *Chirality*, 2010. 22(10): p. 896-900.
26. 赵莉, 高浓度皮质醇介导慢性心理应激和动脉粥样硬化的炎症机制研究——硕士论文. 2009, 山东大学.
27. Jain, N., I. Naseem, and J. Ahmad, Evaluation of DNA damage and metabolic syndrome parameters in diabetic rabbits supplemented with antioxidants. *Fundam Clin Pharmacol*, 2009. 23(2): p. 197-205.
28. Minami, M., et al., High infusion pressure in conjunction with vitreous surgery alters the morphology and function of the retina of rabbits. *Acta Ophthalmol Scand*, 2007. 85(6): p. 633-9.
29. Sugiyama, T., et al., Effect of P2X7 receptor activation on the retinal blood velocity of diabetic rabbits. *Arch Ophthalmol*, 2006. 124(8): p. 1143-9.
30. Okuno, T., et al., Glutamate level in optic nerve head is increased by artificial elevation of intraocular pressure in rabbits. *Exp Eye Res*, 2006. 82(3): p. 465-70.
31. Kobayashi, K., et al., Combination of in vivo angiopoietin-1 gene transfer and autologous bone marrow cell implantation for functional therapeutic angiogenesis. *Arterioscler Thromb Vasc Biol*, 2006. 26(7): p. 1465-72.
32. Kawai, T., et al., Hereditary postprandial hypertriglyceridemic rabbit exhibits insulin resistance and central obesity: a novel model of metabolic syndrome. *Arterioscler Thromb Vasc Biol*, 2006. 26(12): p. 2752-7.
33. Kato, M., et al., Severity of hyperlipidemia does not affect antiatherosclerotic effect of an angiotensin II receptor antagonist in apolipoprotein E-deficient mice. *J Cardiovasc Pharmacol*, 2006. 47(6): p. 764-9.
34. Kato, M., et al., Effect of combined treatment with an angiotensin II receptor antagonist and an HMG-CoA reductase inhibitor on atherosclerosis in genetically hyperlipidemic rabbits. *J Cardiovasc Pharmacol*, 2005. 46(4): p. 556-62.
35. Hirao, M., et al., Effects of adenosine on optic nerve head circulation in rabbits. *Exp Eye Res*, 2004. 79(5): p. 729-35.
36. Goto, W., et al., Amelioration by topical bunazosin hydrochloride of the impairment in ocular blood flow caused by nitric oxide synthase inhibition in rabbits. *J Ocul Pharmacol Ther*, 2003. 19(1): p. 63-73.
37. Thakur, N.K., et al., Anti-atherosclerotic effect of beta-blocker with nitric oxide-releasing action on the severe atherosclerosis. *J Cardiovasc Pharmacol*, 2002. 39(2): p. 298-309.
38. Okuno, T., et al., Evidence that nitric oxide is involved in autoregulation in optic nerve head of rabbits. *Invest Ophthalmol Vis Sci*, 2002. 43(3): p. 784-9.
39. Murakami, S., et al., Taurine suppresses development of atherosclerosis in Watanabe heritable hyperlipidemic (WHHL) rabbits. *Atherosclerosis*, 2002. 163(1): p. 79-87.
40. Kida, T., et al., The mechanism and change in the optic nerve head (ONH) circulation in rabbits after glucose loading. *Curr Eye Res*, 2001. 22(2): p. 95-101.
41. Sugiyama, T., et al., Effect of continuous intravenous infusion of carteolol chloride on tissue blood flow in rabbit optic nerve head. *Jpn J Ophthalmol*, 1999. 43(6): p. 490-4.
42. Li, J., et al., Antiatherogenic effect of angiotensin converting enzyme inhibitor (benazepril) and angiotensin II receptor antagonist (valsartan) in the cholesterol-fed rabbits. *Atherosclerosis*, 1999. 143(2): p. 315-26.
43. Nakamura, M., S. Abe, and M. Tanaka, Oral administration of NO synthase inhibitor failed to promote arteriosclerotic lesions in the aorta and the coronary arteries of rabbits fed cholesterol. *Atherosclerosis*, 1998. 141(1): p. 53-60.
44. Hayashi, T., et al., The beta adrenoreceptor antagonist, nipradilol, preserves the endothelial nitric oxide response in atherosclerotic vessels of rabbit. *Life Sci*, 1997. 61(14): p. 1379-87.